

Exact diagonalization results for an anharmonically trapped Bose-Einstein condensate

S. Bargi, G. M. Kavoulakis, and S. M. Reimann

Mathematical Physics, Lund Institute of Technology, P. O. Box 118, S-22100 Lund, Sweden

(Dated: February 2, 2008)

We consider bosonic atoms that rotate in an anharmonic trapping potential. Using numerical diagonalization of the Hamiltonian, we identify the various phases of the gas as the rotational frequency of the trap and the coupling between the atoms are varied.

PACS numbers: 05.30.Jp, 03.75.Hh, 67.40.Db

I. INTRODUCTION

The behavior of confined, Bose-Einstein condensed atoms under rotation has been studied extensively in recent years, both experimentally [1, 2, 3, 4, 5], as well as theoretically. One factor which is very crucial in these studies is the form of the confinement. In most of the experiments performed on cold gases of atoms, the trapping potential is harmonic. It is interesting, however, that the harmonic trapping is special in many respects, as we show below. For this reason recent theoretical studies [6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21], as well as the experiment of Ref. [22], have considered traps with other functional forms, in which the trapping potential grows more rapidly than quadratically at distances far away from the center of the cloud. Such trapping potentials introduce many novel phases.

Motivated by this observation, we examine the lowest state of a Bose-Einstein condensate in a rotating anharmonic trap. Up to now, all theoretical studies that have been performed on this problem were based upon the mean-field approximation. However, especially in the case of effective attractive interactions between the atoms [23, 24], the use of the mean-field approximation is questionable, as we explain in more detail below. In this study, we present “exact” results – for small numbers of atoms – from numerical diagonalization of the Hamiltonian, which goes beyond the mean-field approximation.

Experimentally it is possible to tune both the frequency of rotation of the trap, as well as the strength of the interatomic interaction. Motivated by these facts, we consider here a fixed trapping potential and examine the various phases and the corresponding phase diagram as function of the rotational frequency and of the coupling.

In what follows we present our model in Sec. II. In Sec. III we discuss the general structure of the phase diagram. In Sec. IV we describe the details of the numerical diagonalization. In Secs. V and VI we present and analyze our results for effective repulsive and attractive interatomic interactions, respectively. Finally, in Sec. VII we summarize our main results.

II. MODEL

The Hamiltonian that we consider consists of the single-particle part $h(\mathbf{r}_i)$, and of the interaction $V(\mathbf{r}_i - \mathbf{r}_j)$, which is taken to be of the form of a contact potential, $V(\mathbf{r}_i - \mathbf{r}_j) = U_0 \delta(\mathbf{r}_i - \mathbf{r}_j)$,

$$H = \sum_{i=1}^N h(\mathbf{r}_i) + \frac{1}{2} \sum_{i \neq j=1}^N V(\mathbf{r}_i - \mathbf{r}_j). \quad (1)$$

Here N is the number of atoms, and $U_0 = 4\pi\hbar^2 a/M$, where a is the scattering length for elastic atom-atom collisions, and M is the atom mass. The single-particle part of the Hamiltonian has the usual form

$$h(\mathbf{r}) = -\frac{\hbar^2}{2M} \nabla^2 + V_{\text{trap}}(\mathbf{r}), \quad (2)$$

where the trap is assumed to be symmetric around the z axis. This is also taken to be the axis of rotation. In cylindrical coordinates ρ, ϕ , and z ,

$$V_{\text{trap}}(\mathbf{r}) = \frac{1}{2} M \omega^2 \rho^2 \left(1 + \lambda \frac{\rho^2}{a_0^2} \right) + V(z). \quad (3)$$

Here ω is the trap frequency of the harmonic part of V_{trap} , λ is a dimensionless constant which is the coefficient of the anharmonic term in V_{trap} (taken to be much smaller than unity in this study, as is also the case in the experiment of Ref. [22]), and $a_0 = \sqrt{\hbar/M\omega}$ is the oscillator length. Along the z axis we assume that the trapping potential $V(z)$ is sufficiently strong that the typical separation between single-particle energy levels is much larger than the typical interaction energy. This assumption implies that the motion along the z axis is frozen out and our problem is essentially two-dimensional.

III. GENERAL STRUCTURE OF THE PHASE DIAGRAM

As mentioned in the introduction, motivated by the experimental situation, we here examine the phase diagram as function of the rotational frequency Ω of the trap and of the strength of the interaction. It is natural to measure Ω in units of ω and the strength of the interaction $\sim nU_0$, where n is the atomic density, in units of

the oscillator energy $\hbar\omega$. The corresponding dimensionless ratio $nU_0/\hbar\omega$ is $\sim \sigma a$, where $\sigma = N/Z$ is the density per unit length. Here Z is the width of the atomic cloud along the z axis.

It is instructive to first consider the general structure of the phase diagram [10, 14, 15]. This consists of several distinct phases. Let us start with repulsive interactions, $a > 0$. For a fixed Ω , as σa increases, the total energy eventually becomes dominated by the interaction energy, which is minimized via the formation of singly quantized vortex states. This is similar to vortex formation in liquid Helium, or in harmonically trapped gases.

On the other hand, for a constant interaction strength, as the rotational frequency increases – and the single-particle part is the dominant part of the Hamiltonian – the total energy of the gas is minimized via formation of multiply quantized vortices. This is most easily seen if one thinks of the effective potential felt by the atoms, which consists of V_{trap} minus the centrifugal potential,

$$\begin{aligned} V_{\text{eff}}(\mathbf{r}) &= V_{\text{trap}}(\mathbf{r}) - \frac{1}{2}M\Omega^2\rho^2 = \\ &= \frac{1}{2}M(\omega^2 - \Omega^2)\rho^2 + \frac{\lambda}{2}M\omega^2\frac{\rho^4}{a_0^2} + V(z). \end{aligned} \quad (4)$$

For the form of V_{trap} we have considered, V_{eff} takes the form of a mexican-hat shape for $\Omega > \omega$, and thus the atoms prefer to reside along the bottom of this potential. This is precisely the density distribution that corresponds to multiply-quantized vortex states. From Eq. (4) it is also clear that for harmonic trapping, $\lambda = 0$, Ω cannot exceed ω , since the atoms will fly apart. For any other potential that grows more rapidly than ρ^2 at large distances, the effective potential is bounded, independently of Ω . Finally, when both σa and Ω/ω are sufficiently large, there is a third, mixed phase consisting of a multiply-quantized vortex state at the center of the cloud, surrounded by singly-quantized vortices. This configuration in a sense compromises between the single-particle energy and the interaction energy.

Turning to the case of effective attractive interactions, $a < 0$, for small enough $\sigma|a|$, the phase diagram is, crudely speaking, symmetric with respect to σa . The lowest state of the gas is determined by the single-particle part of the Hamiltonian, and thus the corresponding state still consists of multiply quantized vortex states. For higher values of $\sigma|a|$, these states are again unstable against the formation of a combination of multiply quantized and singly quantized vortices, as in the mixed phase described for positive σa . The only difference is that the single-particle density distribution resembles that of a localized “blob” rotating around the minimum of the effective potential V_{eff} [16]. For even higher values of $\sigma|a|$, the phase that minimizes the energy is that of the center of mass excitation first seen in harmonically trapped atoms [23, 24]. In this phase the angular momentum is carried by the center of mass. The single-particle density distribution resembles that of the “mixed” phase (i.e., a localized blob), although the structure of the many-body




FIG. 1: Schematic phase diagram which shows the various phases as the rotational frequency and the coupling are varied.

wavefunction is very different. Finally, for sufficiently large values of $\sigma|a|$ the gas collapses [25, 26, 27]. Figure 1 shows a schematic phase diagram.

IV. NUMERICAL DIAGONALIZATION OF THE HAMILTONIAN

As mentioned earlier, in the present study we perform numerical diagonalization of the Hamiltonian for a given atom number N and angular momentum $L\hbar$. To achieve this, we first solve the single-particle eigenvalue problem numerically,

$$\hbar\phi_{n_r,m} = E_{n_r,m}\phi_{n_r,m}, \quad (5)$$

using the Numerov method. The eigenfunctions and eigenenergies are characterized by two quantum numbers, the number of radial nodes n_r , and the number of quanta of angular momentum m . Making the simplified assumption of weak interactions, $\sigma|a| \ll 1$, we perform our calculation within the approximation of the lowest Landau level, $n_r = 0$ and $m \geq 0$. States with $n_r \neq 0$ or negative m are higher in energy by a term of order $\hbar\omega$. This assumption allows us to consider relatively large values of N and L . One should remember that while for repulsive interactions σa can have any value, for attractive interactions, $\sigma|a| \lesssim 1$, as otherwise the system collapses. Therefore, the lowest Landau level approximation is more restrictive for repulsive interactions, yet it gives a rather good description of the various phases, even in this case.

Having found the single-particle eigenstates, we then set up the Fock states, which are the eigenstates of the number operator $\hat{N}$ and of the total angular momentum




FIG. 2: The phase diagram calculated within the mean-field approximation for $\lambda = 0.005$ from Refs. [14, 15, 16]. The “c.m.” phase is the center of mass phase. The stars and the three dashed horizontal lines represent some of the values of σa and Ω/ω that have been examined and analyzed in the present study.

$\hat{L}$, respectively. As a final step we set up the corresponding Hamiltonian matrix between the various Fock states, which we diagonalize numerically. The diagonal matrix elements consist both of the single-particle part, as well as the interaction. The only non-zero off-diagonal matrix elements result from the interaction. The output of the Hamiltonian gives us not only the lowest state, but the whole energy spectrum. Since our calculation is performed for fixed angular momentum, in order to work with a fixed Ω we consider the energy in the rotating frame, which is given by the usual transformation

$$H' = H - L\Omega. \quad (6)$$

Having found the many-body eigenstates and eigenvalues for some range of L , then for a fixed Ω we identify the eigenstate with the lowest eigenenergy according to Eq. (6). The many-body eigenfunction that corresponds to the lowest eigenenergy – which is expressed in terms of the Fock states – is then analyzed in terms of the occupancy of the single particle states. The calculation of the occupancies is a trivial operation and, remarkably, each of the phases described in the previous section is characterized by a distinct distribution. This fact provides indisputable evidence for the phases we expect to get from the arguments of the previous section and from the predictions of mean-field.




FIG. 3: The occupancy of the single-particle states of angular momentum m for $\lambda = 0.005$, $\sigma a = 0.04$, $\Omega/\omega = 1.005$, and for $N = 5$ (higher), $N = 10$ (middle), and $N = 20$ (lower). These graphs demonstrate that the occupancy of all states with $m \neq 1$ is, to leading order, $1/N$. The occupancy of the $m = 1$ state is unity minus corrections of order $1/N$ to leading order. In the mean-field approximation this phase corresponds to a single vortex state.

V. RESULTS – REPULSIVE INTERACTIONS

We turn now to the analysis of our results, starting with the case of repulsive interactions. As we argued earlier, for sufficiently weak interactions we expect the phase of multiple quantization to have the lowest energy. For convenience we compare our results with those of the phase diagram of Ref. [14]. We thus choose $\lambda = 0.005$ [28], and evaluate the occupancy of the single particle states for $\sigma a = 0.04$, $\Omega/\omega = 1.005$, for $N = 5, 10$, and 20 atoms, as shown in Fig. 3. This graph demonstrates clearly that as N increases, the occupancy of the $m = 1$ state approaches unity, while the occupancy of all other states tends to zero (scaling as $1/N$ to leading order). In the mean-field description the order parameter ψ is simply

$$\psi = \phi_{0,1}, \quad (7)$$




FIG. 4: The occupancy of the single-particle states of angular momentum m for $\lambda = 0.005$, $\sigma a = 0.04$, $N = 10$ atoms, and $\Omega/\omega = 1.005$ (top left), $\Omega/\omega = 1.014$ (top right), $\Omega/\omega = 1.020$ (bottom left), $\Omega/\omega = 1.026$ (bottom right). The angular momentum of the gas changes discontinuously between successive values of m as Ω increases. These phases correspond to multiply-quantized vortex states.




FIG. 5: The total angular momentum of the gas in its lowest-energy state as a function of Ω/ω , for fixed $\lambda = 0.005$, $\sigma a = 0.04$, and $N = 5$. The stars denote the result of a perturbative, mean-field calculation from Ref. [15].

which represents a single vortex state located at the center of the cloud.

In Fig. 4 we calculate the occupancies of the single particle states for $\lambda = 0.005$, $\sigma a = 0.04$ and for four values of $\Omega/\omega = 1.005, 1.014, 1.020$, and 1.026 . This graph confirms that indeed the system (in its lowest state) under-




FIG. 6: The occupancy of the single-particle states of angular momentum m for $\lambda = 0.005$, $\sigma a = 0.1$, $\Omega/\omega = 1.0025$ and $N = 10$ (higher), $N = 20$ (lower). Here $L/N = 1.8$. These graphs demonstrate that the dominant states are the ones with $m = 0, 2$, and 4 , in agreement with the mean-field calculations of Refs. [14, 15]. In the mean-field approximation this corresponds to a “mixed” phase discussed in the text.

goes discontinuous transitions between states of multiple quantization of successive values of m . Figure 5 shows the corresponding angular momentum per particle $l = L/N$ in the lowest state of the gas versus Ω/ω , for a fixed value of $\sigma a = 0.04$. This graph consists of discontinuous jumps in l as Ω increases. The asterisks in this graph show the result of mean field [15], which compares well with our result for small Ω . For larger Ω the agreement becomes worse. This is because the contribution of the anharmonic term to the energy was calculated perturbatively in λ in Ref. [15]. This correction increases quadratically with m , and for larger values of this quantum number, the perturbative result deviates from the exact.

For repulsive interactions, as σa increases, mean-field theory predicts that the system undergoes a continuous transition from some multiply-quantized vortex state of angular momentum m_0 to a “mixed” state which, to leading order, is a linear superposition of three states with angular momentum m_1 , m_0 , and m_2 , with $m_1 + m_2 = 2m_0$. Thus the order parameter has the form

$$\psi = c_{m_1} \phi_{0,m_1} + c_{m_0} \phi_{0,m_0} + c_{m_2} \phi_{0,m_2}, \quad (8)$$

where the coefficients are complex numbers. We have confirmed that our method gives a similar transition, as




FIG. 7: The occupancy of the single-particle states of angular momentum m for $\lambda = 0.005$, $\sigma a = -0.001$, $N = 5$ atoms, and $\Omega/\omega = 1.013$ (top left), $\Omega/\omega = 1.018$ (top right), $\Omega/\omega = 1.023$ (bottom left), $\Omega/\omega = 1.028$ (bottom right). In the mean-field approximation this phase corresponds to multiply-quantized vortex states.

shown in Fig. 6. In this figure we plot the occupancy of the single-particle states for $\sigma a = 0.1$, $\Omega/\omega = 1.0025$, and $\lambda = 0.005$, for $N = 10$ and $N = 20$. From these two graphs we see that the states with $m_1 = 0$, $m_0 = 2$, $m_2 = 4$ are the dominant ones for large N . This is again in agreement with the prediction of mean-field. Finally, the angular momentum per particle $l = L/N$ is 1.8 in this state.

VI. RESULTS – ATTRACTIVE INTERACTIONS

For an effective attractive interaction between the atoms the corresponding phase diagram is even richer. We have obtained evidence for all the phases that are expected, namely the multiply-quantized vortex states, the mixed phase that consists of singly and multiply quantized vortices, the one involving center of mass excitation, and finally the unstable one.

Figure 7 shows the occupancies of the single-particle states in the lowest state of the gas for a fixed $\lambda = 0.005$ and $\sigma a = -0.001$, and for four values of $\Omega/\omega = 1.013, 1.018, 1.023$, and 1.028 . As in Fig. 4, as Ω increases, we see that states of multiple quantization $\phi_{0,m}$ with successive values of m become macroscopically oc-




FIG. 8: The angular momentum per particle in the state of lowest energy as a function of Ω/ω , for $\lambda = 0.005$, $N = 5$, and $\sigma a = -0.001$ (higher), and $\sigma a = -0.002$ (lower). The plateaus correspond to multiply-quantized vortex states, which become more narrow as $|\sigma a|$ increases (as also shown in Fig. 2), in agreement with the mean-field approximation [16]. The small wiggles in the curves between the plateaus are an artifact of the discreteness of L that we have considered.

cupied. In Fig. 8 we plot the angular momentum per particle $l = L/N$ versus Ω/ω for $\sigma a = -0.001$ and -0.002 . Here there is a difference as compared to repulsive interactions, which is also consistent with the prediction of mean-field theory. In order for the gas to get from some state of multiple quantization m_0 to $m_0 + 1$ (corresponding to the plateaus in Fig. 8), it has to go through some “mixed state”, and thus these transitions are continuous. Furthermore, the width (in Ω/ω) of the plateaus decreases with increasing $|\sigma a|$. This effect is clearly seen between the two graphs in Fig. 8.

To get evidence for the mixed phase, we increase $|\sigma a|$, $\sigma a = -0.004$, and plot the occupancies of the single particle states in Fig. 9 for a fixed $\Omega/\omega = 1.013$, and $N = 5, 10$, and 20 . Here L/N is 1.0. The dominant single-particle states are the ones with $m = 0, 1$, and 2 , with a very small admixture of $m = 3$. Again, this is in agreement with mean-field theory, which predicts that close to the phase boundary, any multiply-quantized vortex state of strength m_0 is unstable against a state with three components of angular momentum $m_0 - 1, m_0$, and $m_0 + 1$,

$$\psi = c_{m_0-1}\phi_{0,m_0-1} + c_{m_0}\phi_{0,m_0} + c_{m_0+1}\phi_{0,m_0+1}. \quad (9)$$




FIG. 9: The occupancy of the single-particle states of angular momentum m for $\lambda = 0.005$, $\sigma a = -0.004$, $\Omega/\omega = 1.013$, and for $N = 5$ (higher), $N = 10$ (middle), and $N = 20$ (lower). Here $L/N = 1.0$. These graphs demonstrate that the dominant states are the ones with $m = 0, 1$, and 2 , with a small admixture of the state with $m = 3$, in agreement with Ref. [16]. In the mean-field approximation, this corresponds to the “mixed” phase discussed in the text.

In such a linear superposition of single-particle states, the corresponding single-particle density distribution may (and it actually does) look very much like that of a localized blob [16], although this is a phase involving vortex excitation.

The phase of the center of mass excitation expected in this problem [16] was also clearly seen in our calculation for even more negative values of σa . This phase was first discovered in harmonic traps [23, 24]. Remarkably, the many-body wavefunction can be written analytically and has a very different structure as compared to the product form assumed within the mean-field approximation. This fact makes the validity of the mean-field approximation questionable for attractive interactions.

The occupancy of the single-particle states can also be expressed analytically. In Ref. [23] it has been shown that the occupancy $|c_m|^2$ of a single particle state with angular momentum m in a many-body state with center




FIG. 10: The crosses show the occupancy of the single-particle states of angular momentum m for $\lambda = 0.005$, $\sigma a = -0.05$, $N = 5$ atoms, and $\Omega/\omega = 1.012$ (top left), $\Omega/\omega = 1.014$ (top right), $\Omega/\omega = 1.016$ (middle left), $\Omega/\omega = 1.018$ (middle right), $\Omega/\omega = 1.022$ (bottom left), and $\Omega/\omega = 1.026$ (bottom right). This phase has a very large overlap with that of the center of mass excitation. The circles show the results calculated from Eq. (10).

of mass excitation of L units of angular momentum and N atoms is

$$|c_m(L, N)|^2 = \frac{(N-1)^{L-m} L!}{N^L (L-m)! m!}. \quad (10)$$

The crosses in Fig. 10 show the occupancy of the single-particle states calculated numerically for $\sigma a = -0.05$, $\lambda = 0.005$, $N = 10$, and six values of $\Omega/\omega = 1.012, 1.014, 1.016, 1.018, 1.022$, and 1.026 . The circles represent $|c_m|^2$ of Eq. (10). We confirmed that the difference between the occupancies calculated within our study and the prediction of Eq. (10) decreases with increasing N and increasing $\sigma|a|$ (as long as the gas is stable). These data again provide clear evidence for the phase of center of mass excitation.

It should also be mentioned that the phase boundary between the mixed phase and the one of center of mass excitation lies much lower than the one calculated variationally in Ref. [16]. The reason is that the phase boundary of Ref. [16] (for these two specific phases) is just an upper bound, and the present calculation is consistent with this fact.

Finally, the unstable phase [25, 26, 27] also shows up

in our model. This is seen in the present calculation via the sign of the chemical potential which becomes negative for a value of $\sigma a \approx -1$. This result is inconsistent with the assumption of weak interactions, $\sigma|a| \ll 1$, as the approximation of lowest Landau level breaks down. To get a quantitatively accurate description of this instability one would have to include higher Landau levels.

VII. SUMMARY

To summarize, in the present study we considered bosonic atoms that rotate in an anharmonic trap. We attacked this problem with the method of numerical diagonalization of the Hamiltonian, which goes beyond the mean-field approximation but also forced us to consider small numbers of atoms. Apart from the assumption of

weak interatomic interactions, our results are exact. We got evidence for all the phases that are expected, namely vortex excitation of multiple quantization, single quantization, and mixed; center of mass motion, and finally the unstable phase. The location of the phase boundaries were consistent with those calculated within the mean field approximation.

Acknowledgments

We thank Yongle Yu for his assistance on the numerical calculations. This work was partly financed by the European Community project ULTRA-1D (NMP4-CT-2003-505457), the Swedish Research Council, and the Swedish Foundation for Strategic Research.

-
- [1] M. R. Matthews, B. P. Anderson, P. C. Haljan, D. S. Hall, C. E. Wieman, and E. A. Cornell, *Phys. Rev. Lett.* **83**, 2498 (1999).
 - [2] K. W. Madison, F. Chevy, W. Wohlleben, and J. Dalibard, *Phys. Rev. Lett.* **84**, 806 (2000); F. Chevy, K.W. Madison, and J. Dalibard, *Phys. Rev. Lett.* **85**, 2223 (2000); K. W. Madison, F. Chevy, V. Bretin, and J. Dalibard, *Phys. Rev. Lett.* **86**, 4443 (2001).
 - [3] J. R. Abo-Shaer, C. Raman, J. M. Vogels, and W. Ketterle, *Science* **292**, 476 (2001).
 - [4] P. C. Haljan, I. Coddington, P. Engels, and E. A. Cornell, *Phys. Rev. Lett.* **87**, 210403 (2001); P. Engels, I. Coddington, P. C. Haljan, and E. A. Cornell, *Phys. Rev. Lett.* **89**, 100403 (2002).
 - [5] P. Engels, I. Coddington, P. C. Haljan, V. Schweikhard, and E. A. Cornell, *Phys. Rev. Lett.* **90**, 170405 (2003).
 - [6] A. L. Fetter, *Phys. Rev. A* **64**, 063608 (2001).
 - [7] E. Lundh, *Phys. Rev. A* **65**, 043604 (2002).
 - [8] K. Kasamatsu, M. Tsubota, and M. Ueda, *Phys. Rev. A* **66**, 053606 (2002).
 - [9] U. R. Fischer and G. Baym, *Phys. Rev. Lett.* **90**, 140402 (2003).
 - [10] G. M. Kavoulakis and G. Baym, *New Journal of Physics* **5**, 51.1 (2003).
 - [11] T. P. Simula, A. A. Penckwitt, and R. J. Ballagh, *Phys. Rev. Lett.* **92**, 060401 (2004).
 - [12] E. Lundh, A. Collin, and K.-A. Suominen, *Phys. Rev. Lett.* **92**, 070401 (2004).
 - [13] A. Aftalion and I. Danaila, *Phys. Rev. A* **69**, 033608 (2004).
 - [14] A. D. Jackson, G. M. Kavoulakis, and E. Lundh, *Phys. Rev. A* **69**, 053619 (2004).
 - [15] A. D. Jackson and G. M. Kavoulakis, *Phys. Rev. A* **70**, 023601 (2004).
 - [16] G. M. Kavoulakis, A. D. Jackson, and G. Baym, *Phys. Rev. A* **70**, 043603 (2004).
 - [17] E. Lundh, A. Collin, and K.-A. Suominen, *Phys. Rev. A* **71**, 023613 (2005).
 - [18] S. Stringari, A. Fetter, *Phys. Rev. Lett.* **94**, 100402 (2005).
 - [19] H. Fu and E. Zaremba, e-print cond-mat/0508515.
 - [20] N. Papanicolaou, S. Komineas, and N. R. Cooper, e-print cond-mat/0508563.
 - [21] A. Collin, e-print cond-mat/0509256.
 - [22] V. Bretin, S. Stock, Y. Seurin, and J. Dalibard, *Phys. Rev. Lett.* **92**, 050403 (2004).
 - [23] N. K. Wilkin, J. M. F. Gunn, and R. A. Smith, *Phys. Rev. Lett.* **80**, 2265 (1998).
 - [24] B. Mottelson, *Phys. Rev. Lett.* **83**, 2695 (1999).
 - [25] G. Baym and C. J. Pethick, *Phys. Rev. Lett.* **76**, 6 (1996).
 - [26] M. Ueda and A. J. Leggett, *Phys. Rev. Lett.* **80**, 1576 (1998).
 - [27] E. Mueller and G. Baym, *Phys. Rev. A* **62**, 053605 (2000).
 - [28] In Ref. [14] the value of λ that was used was 0.05, however here we use $\lambda = 0.005$. To compare the present results with those of this reference, the axes of the phase diagram of Ref. [14] have been rescaled. This rescaling is allowed because of the universality of the phase diagram in the limit of small λ , as shown in this reference.